

Breed and Composite Selection

Bob Weaver, University of Missouri-Columbia

With more than sixty (60) breeds of beef cattle present in the United States, the question of “Which breed should I choose?” is a difficult question to answer. The top ten breeds in fiscal year 2007 reported registrations accounting for 93% of the pedigreed beef cattle in the U.S. These top ten breeds and their crosses represent the majority of the genetics utilized in commercial beef production, providing a hint at the breeds that possess the most valuable combinations of traits as recognized by beef producers. The breed, composite or combination of breeds employed in a breeding program can have a large impact on the profitability of a commercial beef operation and the value of animals it produces as they move through the beef complex. The breed or biological type of an animal influences economically important production traits including: growth rate, mature size, reproductive efficiency, milk yield, and carcass merit.

Large differences exist today in the relative performance of various breeds for most economically important traits. These breed differences represent a valuable genetic resource for commercial producers to use in structured crossbreeding systems to achieve an optimal combination of traits matching the cowherd to their production environment and to use sire selection to produce market-targeted progeny. As such, the selection of the ‘right’ breed(s) to use in a breeding program is an important decision for commercial beef producers. The determination of the ‘right’ breed(s) to use is highly dependent on a number of characteristics of a farm or ranch such that not every operation should use the same breed or combination of breeds.

Breed and Composite Defined

A common definition of a breed is a genetic strain or type of domestic livestock that has consistent and inherited characteristics such as coat color or pattern, presence or absence of horns, or other qualitative criteria. However, one can also consider performance traits as common characteristics shared by individuals of a breed. In simple terms, these common characteristics are the performance traits that are often associated with a breed as its reputation has grown over time and represent the core traits for which a breed of livestock has been selected for over time. Breeds differ in the level of performance for various traits as a result of different selection goals of their breeders.

A composite is something that is made up of distinct components. In reference to beef cattle, the term composite generally means that the animal is composed of two or more breeds. A composite breed then is a group of animals of similar breed composition. Composites can be thought of as new breeds and managed as such.

Beef Breed and Composite Characterization

A great deal of research has been conducted over the last 30 years at various federal and state experiment stations to characterize beef breeds in the U.S. These studies have been undertaken to examine the genetic merits of various breeds in a wide range of production environments and management systems. During this time, researchers at the U.S. Meat Animal Research Center

(MARC) have conducted the most comprehensive studies of sire breed genetic merit via their long term Germplasm Evaluation (GPE) project. This project evaluated over 30 sire breeds in a common environment and management system. The data summarized by the MARC scientists consisted of records on more than 20,000 animals born between 1978 and 1991, with a re-sampling of the most popular sire breeds in 1999-2000. The various sire breeds evaluated were mated to Angus, Hereford and crossbred cows. Thus, the data reported were for crossbred progeny. During the study, Angus-Hereford crossbred calves were produced in the study as a control for each cycle of the GPE project.

One of the major outcomes of the GPE project was the characterization of sire breeds for a wide variety of economically important

Table 1. Breeds grouped into biological type by four criteria.^{a,b}

Breed Group	Growth Rate and Mature Size	Percent Retail Product	Age at Puberty	Milk Production
Jersey	X	X	X	XXXXX
Longhorn	X	XXX	XXX	XX
Angus	XXX	XX	XX	XXX
Hereford	XXX	XX	XXX	XX
Red Poll	XX	XX	XX	XXX
Devon	XX	XX	XXX	XX
Shorthorn	XXX	XX	XXX	XXX
Galloway	XX	XXX	XXX	XX
South Devon	XXX	XXX	XX	XXX
Tarentaise	XXX	XXX	XX	XXX
Pinzgauer	XXX	XXX	XX	XXX
Brangus	XXX	XX	XXXX	XX
Santa Gertrudis	XXX	XX	XXXX	XX
Sahiwal	XX	XXX	XXXXX	XXX
Brahman	XXX	XXX	XXXXX	XXX
Nellore	XXX	XXX	XXXXX	XXX
Braunvieh	XXXX	XXXX	XX	XXXX
Gelbvieh	XXXX	XXXX	XX	XXXX
Holstein	XXXX	XXXX	XX	XXXXX
Simmental	XXXXX	XXXX	XXX	XXXX
Maine Anjou	XXXXX	XXXX	XXX	XXX
Salers	XXXXX	XXXX	XXX	XXX
Piedmontese	XXX	XXXXX	XX	XX
Limousin	XXX	XXXX	XXXX	X
Charolais	XXXXX	XXXX	XXXX	X
Chianina	XXXXX	XXXX	XXXX	X

^a Adapted from Cundiff et al. 1993.

^b Increasing number of X's indicate relatively higher levels of trait.

traits. Because all of the animals were in a common management system and production environment, the average differences observed in performance were due to genetic differences. Following the analysis of progeny data, the breeds can be divided into groups based on their biological type for four criteria: 1) Growth rate and mature size 2) Lean to fat ratio 3) Age at puberty, and, 4) Milk production. The breeds evaluated at MARC are grouped by biological type in Table 1. British breeds such as Hereford, Angus, Red Angus and Shorthorn are moderate in growth and mature size, relatively higher in carcass fat composition, reach puberty at relatively younger ages and are moderate in milk production. Continental European breeds, with a heritage that includes milk production, including Simmental, Maine-Anjou, and Gelbvieh tend to have high growth rates, larger mature sizes, moderate ages at puberty and relatively high levels of milk production. Another group of Continental European breeds, with a heritage of meat and draft purposes, including Charolais, Chianina and Limousin tend to have high growth rate, large mature size, older ages at puberty, very lean carcasses and low milk production.

Another way to compare the relative genetic merit of breeds for various performance traits is through conversion of their EPD to a common base. This can be accomplished using the across breed EPD adjustments published each year in the proceedings of the Beef Improvement Federation's annual meeting. These adjustments are generated by researchers at MARC. Table 2 lists the across breed adjustment factors that are added to the EPD of an animal of a specified breed to put that animal's EPD on an Angus base (Kuehn and Thallman, 2009). Table 3 presents the average across breed EPD of animals born in 2007 as reported from 2009 genetic evaluations from the most widely used breeds on a common genetic base (Angus). Differences in across breed EPD averages represent genetic differences for each trait. Table 3 provides a more contemporary look at the differences in breed genetic potential for various traits and accounting for genetic trends occurring in each breed due to selection. Due to selection pressure placed on growth and maternal traits over time, many breeds have made considerable gains in those traits. In some cases, the large gains in performance have resulted in subtle changes in the overall biological type of a breed.

Table 2. 2009 adjustment factors to add to EPD of fifteen different breeds to estimate across breed EPDs.

Breed	Birth Wt.	Weaning Wt.	Yearling Wt.	Maternal Milk	Marbling Score	Ribeye Area	Fat Thickness
Angus	0.0	0.0	0.0	0.0	0.00	0.00	0.00
Beefmaster	7.7	44.2	44.0	2.6			
Brahman	11.2	36.3	2.2	29.0			
Brangus	4.7	21.9	19.9	2.4			
Braunvieh	7.5	21.4	12.8	30.6	-0.26	0.78	-0.149
Charolais	9.7	38.2	51.9	5.6	-0.50	0.63	-0.244
Chiangus	4.1	-19.6					
Gelbvieh	4.5	1.7	-12.6	9.9			
Hereford	2.9	-2.8	-16.1	-17.5	-0.36	-0.24	-0.057
Limousin	4.2	-3.4	-28.6	-14.2	-0.80	0.93	
Maine-Anjou	5.5	-10.7	-22.8	-0.8	-0.92	1.07	-0.197
Red Angus	2.9	-5.4	-4.4	-3.0	-0.01	-0.21	-0.045
Salers	3.4	22.7	52.3	13.1	-0.11	0.78	-0.224
Santa Gertrudis	8.1	17.1					
Shorthorn	6.1	19.9	52.8	23.1	0.06	0.12	-0.133
Simmental	5.5	25	22.4	13.7	-0.60	0.92	-0.193
South Devon	4.5	6.9	-1.4	-6.5	-0.32	0.39	-0.131
Tarentaise	2.5	29.7	17.9	22.2			

^a Kuehn and Thallman, 2009.

Table 3. Average Across-Breed EPD for animals born in 2007 by breed from 2009 genetic evaluations and 2009 USDA-MARC Across-Breed EPD adjustment factors.

Breed	Birth Wt.	Weaning Wt.	Yearling Wt.	Maternal Milk	Marbling Score	Ribeye Area	Fat Thickness
Angus	2.2	43.5	80.0	20.5	0.31	0.15	0.01
Beefmaster	8.2	51.5	56.5	4.6			
Brahman	13.0	49.9	24.4	34.8			
Brangus	5.3	43.8	60.1	9.7			
Braunvieh	7.3	22.3	14.3	30.9	-0.25	0.79	-0.15
Charolais	10.3	61.5	93.1	12.1	-0.47	0.80	-0.24
Chiangus	5.3	24.6					
Gelbvieh	5.8	42.7	61.4	27.9			
Hereford	6.4	38.2	51.9	-1.5	-0.33	-0.07	-0.06
Limousin	5.9	39.2	50.5	7.1	-0.80	1.33	
Maine-Anjou	7.4	29.4	56.3	19.2	-0.71	1.23	-0.20
Red Angus	3.2	25.8	50.5	13.1	0.05	-0.16	-0.05
Salers	4.3	40.5	81.9	21.5	-0.11	0.80	-0.22
Santa Gertrudis	8.6	21.1					
Shorthorn	8.3	34.2	76.2	25.6	0.06	0.10	-0.13
Simmental	6.8	57.4	79.9	17.9	-0.47	1.00	-0.18
South Devon	7.1	46.6	74.1	14.9	-0.06	0.49	-0.12
Tarentaise	4.0	33.7	28.9	23.2			

^a Adapted from Kuehn and Thallman, 2009 and Kuehn et al., 2009

Use of Breeds and Composites for Genetic Improvement

Inclusion or exclusion of germplasm from a breed (or composite) is a valuable selection tool for making rapid directional changes in genetic merit for a wide range of traits. Changes in progeny phenotype that occur when breeds are substituted in a breeding program come from two genetic sources.

The first source of genetic impact from a substitution of a breed comes through changes in the additive genetic effects or breeding values that subsequent progeny inherit from their sire and dam. Additive genetic merit is the portion of total genetic merit that is transmissible from parent to offspring and on which traditional selection decisions are made. In other words, additive genetic effects are heritable. EPD are estimates of one-half of the additive genetic merit. The difference in average performance for a trait observed between two breeds is due primarily to differences in additive genetic merit.

The second source of genetic change is due to non-additive genetic effects. Non-additive effects include both dominance and epistatic effects. Dominance effects arise from the interactions of paired genes at each locus. Epistatic effects are the interaction of genes across loci. The sum of these two interactions result in heterosis observed in crossbred animals. Since each parent only contributes one gene to an offspring and dominance effects depend on the interaction of a pair of genes, a parent cannot transmit dominance effects to its progeny within a breed. However, the selection of which breeds and how much of each breed to incorporate into progeny has a large impact on dominance (or heterosis) effects which affect phenotype. Because epistatic effects arise from the interaction of genes at different loci, independent segregation of chromosomes in the formation of gametes causes pairings of genes not to always stay together from one generation to the next. Like dominance effects, epistatic effects are not impacted by mate selection but by the frequency of different alleles and their dominance effects across breeds.

Both additive and non-additive genetic effects can have a significant impact on a particular phenotype; therefore, it is important that both are considered during breed selection. Due to their different modes of inheritance, different tactics must be employed to capture the benefits of each.

Additive genetic merit may be selected for in two distinct ways. First, by the selection of individuals *within* a breed that have superior genetic merit for the trait under selection. Typically this is achieved through the use of EPD to identify selection candidates, although it can also be done through selection for specific alleles using DNA markers. The rate of improvement in phenotypes due to selection within breed is limited by the heritability of the trait. Heritability describes the proportion of phenotypic variation that is controlled by additive genetic variation. So, for traits with moderate to high heritability, considerable progress in progeny phenotype may be achieved through selection of superior animals within the breed as parent stock. The second approach to change additive genetic merit is through the selection of animals from a different breed(s) that excels in the trait under selection. *Across* breed selection can provide rapid change in progeny phenotype given that large differences exist between breeds in a number of economically relevant traits. Selection of superior parent stock from a different breed that excels in a trait is often more effective than selection within a breed (Gregory et al., 1999) as the breed differences have a heritability of nearly 100%.

The use of breed differences to achieve the best overall results across multiple traits may be achieved through the implementation of the concept of breed complementarity. Breeds are complementary to each other when they excel in different traits and their crossbred progeny have desirable levels of performance in a larger number of traits than either of the parent breeds alone. Making breed and mating selections that utilize breed

complementarity provide an effective way to aggregate the core competencies of two or more breeds in the progeny. Moreover, use of breed complementarity can be a powerful strategy to genetically match cows to their production environment and progeny to the market place. For example, a crossbreeding system that mates Charolais bulls to Hereford-Angus crossbred cows utilizes breed complementarity. The Charolais bull contributes growth and carcass yield to progeny genetics while the Hereford-Angus crossbred cows have many desirable maternal attributes and contribute genetics for carcass quality. When considering crossbreeding from the standpoint of producing replacement females, one could select breeds that have complementary maternal traits such that females are most ideally matched to their production environment. Matings to produce calves for market should focus on complementing traits of the cows and fine tuning calf performance (growth and carcass traits) to the market place.

There is an abundance of research that describes the core competencies (biological type) of many of today's commonly used beef breeds as described earlier and listed in Table 1. Traits are typically combined into groupings such as maternal/reproduction, growth and carcass. When selecting animals for a crossbreeding system, breed should be the primary consideration. Breeds selected for inclusion in a mating program will be dependant on a number of factors including current cow herd breed composition, forage and production environment, replacement female development system, and calf marketing endpoint. All of these factors help determine the relative importance of traits for each production phase.

One of the challenges of breed selection is the interaction of the animal's genotype with its production environment. Table 4 describes common production environments by level of feed availability and environmental stress and lists optimal levels of a variety of performance traits (Bullock et al., 2002). Here, feed availability refers to the regular availability of grazed or harvested forage and its quantity and quality. Environmental stress includes parasites, disease, heat and humidity. Ranges for mature cow size are low (800 to 1,000 lb.), medium (1000 to 1,200 lb.), and high (1,200 to 1,400 lb.) Clearly, breed choices should be influenced by the production environment in which they are expected to perform.

Crossing of breeds or lines is the primary method to exploit beneficial non-additive effects called heterosis. Heterosis refers to the superiority of the crossbred animal relative to the average of its straightbred parents and heterosis results from an increase in heterozygosity of a crossbred animal's genetic makeup. Heterozygosity refers to a state where an animal has two different forms of a gene. It is believed that heterosis is primarily the result of gene dominance and the recovery from accumulated inbreeding depression of pure breeds. Heterosis is, therefore, dependant on crossbred animals having a greater percentage of heterozygous animals than is present in straightbred animals. The level of heterozygosity an animal has depends on the random inheritance of copies of genes from its parents. In general, animals that are crosses of unrelated breeds, such as Angus and Brahman, exhibit higher levels of heterosis due to more heterozygosity, than do crosses of more genetically similar breeds such as a cross of Angus and Hereford.

Generally, heterosis generates the largest improvement in lowly heritable traits. Moderate improvements due to heterosis are seen in moderately heritable traits. Little or no heterosis is observed in highly heritable traits. Traits such as reproduction and longevity have low heritability. These traits respond very slowly to

selection since a large portion of the variation observed in them is due to environmental effects and non-additive genetic effects, and a small percentage is due to additive genetic differences. But, heterosis generated through crossbreeding can significantly improve an animal's performance for lowly heritable traits, thus the importance of considering both additive and non-additive genetics when designing mating programs. Crossbreeding has been shown to be an efficient method to improve reproductive efficiency and pre-weaning productivity in beef cattle.

Improvements in cow-calf production due to heterosis are attributable to having both a crossbred cow (called maternal or dam heterosis) and a crossbred calf (called individual or calf heterosis). Differing levels of heterosis are generated when various breeds are crossed. Similar levels of heterosis are observed when members of the *Bos taurus* species, including the British (e.g. Angus, Hereford, Shorthorn) and Continental European breeds (e.g. Charolais, Gelbvieh, Limousin, Maine-Anjou, Simmental), are crossed. Much more heterosis is observed when *Bos indicus*, or Zebu, breeds like Brahman, Nelore and Gir, are crossed with *Bos taurus* breeds. The increase in heterosis observed in British by *Bos indicus* crosses for a trait is usually 2-3 times as large as the heterosis for the same trait observed in *Bos taurus* crossbreds (Koger, 1980). The large increase is especially true with heterosis observed in the crossbred cow. The increase in heterosis is sensible as the there are more genetic differences between species than within a species. Heterosis effects reported in the following tables will be divided and noted into those observed in *Bos taurus* crosses or *Bos taurus* by *Bos indicus* crosses. Table 5 details the individual (crossbred calf) heterosis and Table 6 describes the maternal (crossbred cow) heterosis observed for various important production traits in *Bos taurus* crossbreds. These heterosis estimates are adapted from a report by Cundiff and Gregory, 1999, and summarize crossbreeding experiments conducted in the Southeastern and Midwest areas of the US. Table 7 describes the expected individual heterosis of *Bos taurus* by *Bos indicus* crossbred calves, while Table 8 details the estimated maternal (dam) heterotic effects observed in *Bos taurus* by *Bos indicus* crossbred cows. *Bos taurus* by *Bos indicus* heterosis estimates were derived from breeding experiments conducted in the southern US.

The heterosis adjustments utilized by multi-breed genetic evaluation systems are another example of estimates for individual (due to a calf) and maternal (due to a crossbred dam) heterosis. These heterosis adjustments are present in Table 9 and illustrate the differences in expected heterosis for various breed-group crosses. In general the Zebu (*Bos indicus*) crosses have higher levels of heterosis than the British-British, British-Continental, or Continental-Continental crosses.

The production of crossbred calves yields advantages in both heterosis and the blending of desirable traits from two or more breeds. However, the largest economic benefit of crossbreeding to commercial producers comes from the crossbred cow. Dam heterosis improves both the environment a cow provides for her calf as well as improves her longevity and durability. The improvement of the maternal environment a cow provides for her calf is manifested in improvements in calf survivability to weaning and increased weaning weight. Crossbred cows exhibit improvements in calving rate of nearly 4% and an increase in longevity of more than one year due to heterotic effects. Heterosis results in increases in lifetime productivity of approximately one calf and 600 pounds of calf weaning weight over the lifetime of the cow.

Table 4. Matching genetic potential for different traits to production environments.¹

Production Environment		Traits					
Feed Availability	Stress ²	Milk Production	Mature Size	Ability to Store Energy ³	Resistance to Stress ⁴	Calving Ease	Lean Yield
High	Low	M to H	M to H	L to M	M	M to H	H
	High	M	L to H	L to H	H	H	M to H
Medium	Low	M to H	M	M to H	M	M to H	M to H
	High	L to M	M	M to H	H	H	H
Low	Low	L to M	L to M	H	M	M to H	M
	High	L to M	L to M	H	H	H	L to M
Breed role in terminal crossbreeding systems							
Maternal		M to H	L to H	M to H	M to H	H	L to M
Paternal		L to M	H	L	M to H	M	H

L = Low; M = Medium; H = High.

¹ Adapted from Bullock et al., 2002.

² Heat, cold, parasites, disease, mud, altitude, etc.

³ Ability to store fat and regulate energy requirements with changing (seasonal) availability of feed.

⁴ Physiological tolerance to heat, cold, internal and external parasites, disease, mud, and other factors.

Table 5. Units and percentage of heterosis by trait for *Bos taurus* crossbred calves.

Trait	Heterosis	
	Units	Percentage (%)
Calving Rate, %	3.2	4.4
Survival to Weaning, %	1.4	1.9
Birth Weight, lb	1.7	2.4
Weaning Weight, lb	16.3	3.9
Yearling Weight, lb	29.1	3.8
Average Daily Gain, lb/d	0.08	2.6

Table 6. Units and percentage of heterosis by trait for *Bos taurus* crossbred dams.

Trait	Heterosis	
	Units	Percentage (%)
Calving Rate, %	3.5	3.7
Survival to Weaning, %	0.8	1.5
Birth Weight, lb	1.6	1.8
Weaning Weight, lb	18.0	3.9
Longevity, years	1.36	16.2
Lifetime Productivity		
Number of Calves	.97	17.0
Cumulative Weaning Weight, lb	600	25.3

Crossbreeding can have positive effects on a ranch's bottom line by not only increasing the quality and gross pay weight of calves produced but also by increasing the durability and productivity of the cow factory.

The effects of dam heterosis on the economic measures of cow-calf production have been shown to be very positive. The added value of maternal heterosis ranges from approximately \$50/cow/year to nearly \$100/cow/year depending on the amount of maternal heterosis retained in the cowherd (Ritchie, 1998). Heterosis expressed by dams accounted for an increase in net profit per cow of nearly \$75/cow/year (Davis et al., 1994). Their results suggested that the benefits of dam heterosis on profit were primarily the reduced cost per cow exposed. Crossbred cows had higher reproductive rates, longer productive lives, and required fewer replacements than straightbred cows in their study. All of these factors contribute to reduced cost per cow exposed. Further, they found increased outputs, including growth and milk yield, were offset by increased costs.

When it comes to crossing breeds with the goal of producing high levels of maternal or individual heterosis, not all breeds are equal. Heterosis depends on an animal having two different alleles or alternate forms of a gene at a locus. The likelihood of having different copies of genes at a locus is greater in breeds that are less related than when the breeds crossed are closely related. For instance Angus and Hereford, both British breeds, are more similar than Angus and Simmental (a Continental European breed) which are more similar than Angus (a *Bos taurus* breed) and Brahman (a *Bos indicus* breed). Since heterosis offers considerable advantages to commercial producers in terms of reproductive efficiency, productivity and economic returns, care should be given when selecting breeds for inclusion in a crossbreeding system. Just as breeds differ in the amount of heterosis generated when crossed, crossbreeding systems achieve differing levels of heterosis depending on the number of breeds and their fractions represented in each animal. A more complete discussion on crossbreeding and crossbreeding systems appears in a separate chapter in this manual.

When comparing two breeds for inclusion in a crossbreeding system that offer similar strengths, select the breed that offers the most heterosis when mated to animals of other breed(s) in your system. Table 10 provides estimates of the percentage increase in pairs of alleles at a locus that are different (heterozygosity) when various purebreds are crossed to form F1 progeny. These estimates were developed using the input data and procedures suggested by Roughsedge et al., 2001. It is easy to see that not all breeds offer the same increase in heterozygosity, and therefore, heterosis when crossed. Expected percent heterosis for cow fertility, birth weight,

Table 7. Units and percentage of heterosis by trait for *Bos Taurus* by *Bos indicus* crossbred calves.¹

Trait	Heterosis Units
Calving Rate, % ¹	4.3
Calving Assistance, % ¹	4.9
Calf Survival, % ¹	-1.4
Weaning Rate, % ¹	1.8
Birth Weight, lb ¹	11.4
Weaning Weight, lb ¹	78.5

¹ Adapted from Franke et al. 2005; numeric average of Angus-Brahman, Brahman-Charolais, and Brahman-Hereford heterosis estimates.

Table 8. Units and percentage of heterosis by trait for *Bos Taurus* by *Bos indicus* crossbred dams.^{1,2}

Trait	Heterosis	
	Units	Percentage (%)
Calving Rate, % ¹	15.4	--
Calving Assistance Rate, % ¹	-6.6	--
Calf Survival, % ¹	8.2	--
Weaning Rate, % ¹	20.8	--
Birth Weight, lb ¹	-2.4	--
Weaning Weight, lb ¹	3.2	--
Weaning Wt. per Cow Exposed, lb ²	91.7	31.6

¹ Adapted from Franke et al. 2005; numeric average of Angus-Brahman, Brahman-Charolais, and Brahman-Hereford heterosis estimates.

² Adapted from Franke et al. 2001.

Table 9. Individual (calf) and maternal (dam) heterosis adjustments for British, Continental European, and Zebu breed groups for birth weight, weaning weight and post weaning gain.

Breed Combinations	Birth Weight (lb)		Weaning Weight (lb)		Postweaning Gain (lb)
	Calf	Dam	Calf	Dam	Calf
	Heterosis	Heterosis	Heterosis	Heterosis	Heterosis
British x British	1.9	1.0	21.3	18.8	9.4
British x Continental	1.9	1.0	21.3	18.8	9.4
British x Zebu	7.5	2.1	48.0	53.2	28.2
Continental x Continental	1.9	1.0	21.3	18.8	9.4
Continental x Zebu	7.5	2.1	48.0	53.2	28.2

(Wade Shafer, Am. Simmental Association, personal communication)

survival to weaning and weaning weight was computed according to the procedure outlined by Roughsedge et al., 2001. Table 11 provides the expected heterosis percentage for cow fertility observed in F1 females. Similarly, Tables 12, 13 and 14 provide the expected heterosis percentage for birth weight, survival to weaning and weaning weight, respectively. Note that this study provided no estimates of heterosis for *Bos indicus* breeds such as Brahman, Nelore or Gir as only *Bos taurus* breeds common in the United Kingdom and continental Europe were sampled for biologic material.

Summary

Selection of appropriate breeds for a particular production system can be a challenging task. Consideration during the selection process should be given to a number of criteria (Greiner, 2002) including:

- Climate (frost-free days, growing season, precipitation).
- Quantity, quality and cost of feedstuffs available.
- Production system (availability of labor and equipment).
- Market end points and demands.
- Breed complementarity.
- Cost and availability of seedstock.

The selection of breeds and the genetics they contribute to the cowherd can have a large impact on profitability through the aggregate effects on each of the above criteria. Clearly, breeds need to be selected to fit a specific production system, whether that is selling replacement females, weaned feeder calves, or carcass

Table 10. Increase in heterozygosity of F1 animals when respective breeds are crossed.^a

Breed	A	C	Ch	G	H	PH	L	MA	Sa	Sh	S	SD
Angus (A)	0.000	0.110	0.193	0.116	0.136	0.110	0.103	0.061	0.151	0.057	0.071	0.088
Charolais (C)	0.110	0.000	0.134	0.093	0.148	0.141	0.050	0.096	0.048	0.096	0.059	0.148
Chianina (Ch)	0.193	0.134	0.000	0.128	0.262	0.268	0.139	0.165	0.160	0.183	0.162	0.238
Gelbvieh (G)	0.116	0.093	0.128	0.000	0.183	0.189	0.110	0.151	0.114	0.137	0.063	0.149
Hereford (H)	0.136	0.148	0.262	0.183	0.000	0.011	0.172	0.163	0.195	0.110	0.151	0.183
Polled Hereford (PH)	0.110	0.141	0.268	0.189	0.011	0.000	0.166	0.139	0.198	0.089	0.148	0.172
Limousin (L)	0.103	0.050	0.139	0.000	0.172	0.166	0.000	0.081	0.057	0.094	0.071	0.112
Maine-Anjou (MA)	0.061	0.096	0.165	0.151	0.163	0.139	0.081	0.000	0.151	0.057	0.104	0.116
Salers (Sa)	0.151	0.048	0.160	0.114	0.195	0.198	0.057	0.151	0.000	0.175	0.069	0.211
Shorthorn (Sh)	0.057	0.096	0.183	0.137	0.110	0.089	0.094	0.057	0.175	0.000	0.115	0.093
Simmental (S)	0.071	0.059	0.162	0.063	0.151	0.148	0.071	0.104	0.069	0.115	0.000	0.139
South Devon (SD)	0.088	0.148	0.238	0.149	0.183	0.172	0.112	0.116	0.211	0.093	0.139	0.000

^a Adapted from Roughsedge et al., 2001.

Table 11. Cow fertility expected heterosis (%) for F1's (first cross).

Breed	A	C	Ch	G	H	PH	L	MA	Sa	Sh	S	SD
Angus (A)	0.00	7.32	12.87	7.76	9.05	7.32	6.87	4.05	10.04	3.77	4.77	5.85
Charolais (C)	7.32	0.00	8.97	6.21	9.89	9.43	3.35	6.43	3.21	6.43	3.91	9.89
Chianina (Ch)	12.87	8.97	0.00	8.51	17.50	17.85	9.27	10.97	10.66	12.23	10.82	15.90
Gelbvieh (G)	7.76	6.21	8.51	0.00	12.23	12.63	7.32	10.04	7.61	9.12	4.20	9.96
Hereford (H)	9.05	9.89	17.50	12.23	0.00	0.74	11.44	10.89	13.03	7.32	10.04	12.23
Polled Hereford (PH)	7.32	9.43	17.85	12.63	0.74	0.00	11.05	9.27	13.19	5.92	9.89	11.44
Limousin (L)	6.87	3.35	9.27	0.00	11.44	11.05	0.00	5.41	3.77	6.29	4.77	7.47
Maine-Anjou (MA)	4.05	6.43	10.97	10.04	10.89	9.27	5.41	0.00	10.04	3.77	6.95	7.76
Salers (Sa)	10.04	3.21	10.66	7.61	13.03	13.19	3.77	10.04	0.00	11.68	4.62	14.08
Shorthorn (Sh)	3.77	6.43	12.23	9.12	7.32	5.92	6.29	3.77	11.68	0.00	7.69	6.21
Simmental (S)	4.77	3.91	10.82	4.20	10.04	9.89	4.77	6.95	4.62	7.69	0.00	9.27
South Devon (SD)	5.85	9.89	15.90	9.96	12.23	11.44	7.47	7.76	14.08	6.21	9.27	0.00

Table 12. Birth weight expected heterosis (%) for F1's.

Breed	A	C	Ch	G	H	PH	L	MA	Sa	Sh	S	SD
Angus (A)	0.00	2.64	4.65	2.81	3.27	2.64	2.48	1.47	3.63	1.36	1.72	2.11
Charolais (C)	2.64	0.00	3.24	2.24	3.57	3.41	1.21	2.32	1.16	2.32	1.41	3.57
Chianina (Ch)	4.65	3.24	0.00	3.08	6.32	6.45	3.35	3.96	3.85	4.42	3.91	5.75
Gelbvieh (G)	2.81	2.24	3.08	0.00	4.42	4.56	2.64	3.63	2.75	3.30	1.52	3.60
Hereford (H)	3.27	3.57	6.32	4.42	0.00	0.27	4.13	3.94	4.71	2.64	3.63	4.42
Polled Hereford (PH)	2.64	3.41	6.45	4.56	0.27	0.00	3.99	3.35	4.77	2.14	3.57	4.13
Limousin (L)	2.48	1.21	3.35	0.00	4.13	3.99	0.00	1.96	1.36	2.27	1.72	2.70
Maine-Anjou (MA)	1.47	2.32	3.96	3.63	3.94	3.35	1.96	0.00	3.63	1.36	2.51	2.81
Salers (Sa)	3.63	1.16	3.85	2.75	4.71	4.77	1.36	3.63	0.00	4.22	1.67	5.09
Shorthorn (Sh)	1.36	2.32	4.42	3.30	2.64	2.14	2.27	1.36	4.22	0.00	2.78	2.24
Simmental (S)	1.72	1.41	3.91	1.52	3.63	3.57	1.72	2.51	1.67	2.78	0.00	3.35
South Devon (SD)	2.11	3.57	5.75	3.60	4.42	4.13	2.70	2.81	5.09	2.24	3.35	0.00

components. For most producers, that production system should employ a structured crossbreeding system that utilizes two or more breeds. The breeds (and/or composites) chosen should produce calves that are appropriate for the market targeted. Moreover, the system and breeds included should provide a mechanism for the use of crossbred cows that are matched to the production environment in terms of mature size and lactation potential so as to capture the benefits of maternal heterosis. Selection of breeds

that are too large and/or produce too much milk for the forage environment in which they are expected to produce may result in lower reproductive efficiency and increased supplemental feed costs. Selection of breeds provides an opportunity for the beef producer to impact both additive and non-additive genetics of the cowherd. Optimization of these two genetic components requires a disciplined approach to breed selection.

Table 13. Survival to weaning expected heterosis (%) for F1's.

Breed	A	C	Ch	G	H	PH	L	MA	Sa	Sh	S	SD
Angus (A)	0.00	1.90	3.34	2.01	2.35	1.90	1.78	1.05	2.60	0.98	1.24	1.52
Charolais (C)	1.90	0.00	2.33	1.61	2.56	2.44	0.87	1.67	0.83	1.67	1.02	2.56
Chianina (Ch)	3.34	2.33	0.00	2.21	4.54	4.63	2.41	2.85	2.77	3.17	2.81	4.12
Gelbvieh (G)	2.01	1.61	2.21	0.00	3.17	3.28	1.90	2.60	1.98	2.37	1.09	2.58
Hereford (H)	2.35	2.56	4.54	3.17	0.00	0.19	2.97	2.83	3.38	1.90	2.60	3.17
Polled Hereford (PH)	1.90	2.44	4.63	3.28	0.19	0.00	2.87	2.41	3.42	1.54	2.56	2.97
Limousin (L)	1.78	0.87	2.41	0.00	2.97	2.87	0.00	1.40	0.98	1.63	1.24	1.94
Maine-Anjou (MA)	1.05	1.67	2.85	2.60	2.83	2.41	1.40	0.00	2.60	0.98	1.80	2.01
Salers (Sa)	2.60	0.83	2.77	1.98	3.38	3.42	0.98	2.60	0.00	3.03	1.20	3.65
Shorthorn (Sh)	0.98	1.67	3.17	2.37	1.90	1.54	1.63	0.98	3.03	0.00	1.99	1.61
Simmental (S)	1.24	1.02	2.81	1.09	2.60	2.56	1.24	1.80	1.20	1.99	0.00	2.41
South Devon (SD)	1.52	2.56	4.12	2.58	3.17	2.97	1.94	2.01	3.65	1.61	2.41	0.00

Table 14. Weaning weight expected heterosis (%) for F1's.

Breed	A	C	Ch	G	H	PH	L	MA	Sa	Sh	S	SD
Angus (A)	0.00	1.94	3.42	2.06	2.40	1.94	1.82	1.08	2.66	1.00	1.26	1.55
Charolais (C)	1.94	0.00	2.38	1.65	2.62	2.50	0.89	1.71	0.85	1.71	1.04	2.62
Chianina (Ch)	3.42	2.38	0.00	2.26	4.65	4.74	2.46	2.91	2.83	3.25	2.87	4.22
Gelbvieh (G)	2.06	1.65	2.26	0.00	3.25	3.35	1.94	2.66	2.02	2.42	1.11	2.64
Hereford (H)	2.40	2.62	4.65	3.25	0.00	0.20	3.04	2.89	3.46	1.94	2.66	3.25
Polled Hereford (PH)	1.94	2.50	4.74	3.35	0.20	0.00	2.93	2.46	3.50	1.57	2.62	3.04
Limousin (L)	1.82	0.89	2.46	0.00	3.04	2.93	0.00	1.44	1.00	1.67	1.26	1.98
Maine-Anjou (MA)	1.08	1.71	2.91	2.66	2.89	2.46	1.44	0.00	2.66	1.00	1.84	2.06
Salers (Sa)	2.66	0.85	2.83	2.02	3.46	3.50	1.00	2.66	0.00	3.10	1.23	3.74
Shorthorn (Sh)	1.00	1.71	3.25	2.42	1.94	1.57	1.67	1.00	3.10	0.00	2.04	1.65
Simmental (S)	1.26	1.04	2.87	1.11	2.66	2.62	1.26	1.84	1.23	2.04	0.00	2.46
South Devon (SD)	1.55	2.62	4.22	2.64	3.25	3.04	1.98	2.06	3.74	1.65	2.46	0.00

Literature Cited

- Bullock, D., M. Enns, L. Gould, M. MacNeil, and G.P. Rupp. Utilization. 2002. Chapter 6. IN: Guidelines for Uniform Beef Improvement Programs. 8th edition.
- Cundiff, L. V., and K. E. Gregory. 1999. What is systematic crossbreeding? Paper presented at Cattlemen's College, 1999 Cattle Industry Annual Meeting and Trade Show, National Cattlemen's Beef Association. Charlotte, North Carolina, February 11, 1999.
- Cundiff, L. V., and L. D. Van Vleck. 2006. Mean EPDs Reported by Different Breeds. Proc. 38th Annual Research Symposium and Annual Meeting, Beef Improvement Federation, Choctaw, Mississippi. pp 61-66.
- Gregory, K. E., L. V. Cundiff, L. D. Van Vleck. 1999. Composite breeds to use heterosis and breed differences to improve efficiency of beef production. Technical Bulletin Number 1875. ARS-USDA. Washington, DC.
- Cundiff, L. V., F. Szabo, K. E. Gregory, R. M. Koch, M. E. Dikeman, and J. D. Crouse. 1993. Breed comparisons in the Germplasm Evaluation Program at MARC. Pages 124-136 in Proc. 25th Annual Research Symposium and Annual Meeting, Beef Improvement Federation, Asheville, North Carolina.
- Davis, K. C., M. W. Tess, D. D. Kress, D. E. Doornbos, and D. C. Anderson. 1994. Life Cycle Evaluation of Five Biological Types of Beef Cattle in a Cow-Calf Range Production System: II. Biological and Economic Performance. J. Anim. Sci. 72:2591-2598.
- Franke, D. E., S. M. DeRouen, A. R. Williams, and W. E. Wyatt. 2005. Direct and maternal breed additive and heterosis genetic effects for reproductive, preweaning, and carcass traits. Pages 204-209 in Proc. of Symposium on Tropically Adapted Breeds, Regional Project S-1013, American Society of Animal Science, Southern Section Meeting, Little Rock, Arkansas.
- Franke, D. E., O. Habet, L. C. Tawah, A. R. Williams, and S. M. DeRouen. 2001. Direct and maternal genetic effects on birth and weaning traits in multibreed cattle data and predicted performance of breed crosses. J Anim. Sci. 79: 1713-1722.
- Greiner, S. P. 2002. Beef cattle breeds and biological types. Virginia Cooperative Extension Publication 400-803. Virginia Polytechnic Institute and State University. Blacksburg.
- Kuehn, L. A. and R. M. Thallman. 2009. Across-Breed EPD Table. http://www.ars.usda.gov/sp2UserFiles/Place/54380000/GPE/AB_EPD2009News.pdf Accessed on August 19, 2009.
- Kuehn, L.A., L. D. Van Vleck, R. M. Thallman, L. V. Cundiff. 2009. Across-Breed EPD Tables for the Year 2009 Adjusted to Breed Differences for Birth Year of 2007. Proc., Beef Improvement Federation 41st Research Symposium and Annual Meeting, Sacramento, CA. April 30-May 3, 2009. pp. 160-183.
- Ritchie, H. D., 1998. Role of Composites in Future Beef Production Systems. <http://www.msu.edu/~ritchieh/papers/BEEF201.ppt>. Accessed October 2, 2005.
- Roughsedge, T., R. Thompson, B. Villanueva, and G. Simm. 2001. Synthesis of direct and maternal genetic components of economically important traits from beef breed-cross evaluations. J. Anim. Sci. 79:2307-2319